

## 中国外来入侵植物的学名考证\*

闫小玲\*\*, 马金双\*\*

(中国科学院上海辰山植物科学研究中心, 上海 201602)

**摘要:** 基于中国外来入侵植物的相关报道和文献, 发现一些入侵植物的学名使用比较混乱, 一个物种存在两个或两个以上的名称。依据最新植物志资料, 对国内 30 种外来入侵植物学名进行了考证, 包括中名、基名 (若存在)、异名、原产地以及在中国的分布等。

**关键词:** 入侵植物; 中国; 学名; 分类修订

中图分类号: Q 949

文献标识码: A

文章编号: 2095-0845(2011)01-132-11

## Nomenclatural Notes on Alien Invasive Vascular Plants in China

Xiao-Ling YAN\*\*, Jin-Shuang MA\*\*

(Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai 201602, China)

**Abstract:** Based on literatures of invasive plants, the scientific names of thirty alien invasive plants in China are reviewed here, since more than one name were used for one species by different authors. The data from the floras updated were employed to compile the scientific name for each species. The vernacular name, basionym (if present), synonym (s) for each species, along with their distributions in China were provided.

**Key words:** Invasive plants; China; Scientific name; Taxonomic notes

1. *Ageratina adenophora* (Sprengel) R. M. King & H. Robinson, Phytologia **19**: 211. 1970<sup>①</sup>; G. L. Nesom in Fl. N. Amer. **21**: 553. 2006. Basionym *Eupatorium adenophorum* Sprengel, Syst. Veg. **3**: 420. 1826 (Asteraceae).

Sticky snakeroot (破坏草<sup>②</sup>), a native species of Mexico, introduced and naturalized in pantropical and pansubtropical as invasive in Canary Islands, Cambodia, India, Laos, Myanmar, Nepal, Indonesia, Philippines, Thailand, Vietnam, South Africa, Americas, probably all over SE Asia, Australia, Pacific islands, South China Sea Islands, and has been

introduced and naturalized in China since the middle of the nineteenth century: Chongqing, Guangxi, Guizhou, Hebei, Jiangxi, Sichuan, Taiwan and Yunnan. Here *Ageratina* is a restricted circumscription in a broad *Eupatorium*. It is commonly accepted as *E. adenophorum* in China (Li and Xie, 2002; Shi *et al.*, 2004; Guan *et al.*, 2006; Huang *et al.*, 2006; Xu *et al.*, 2006; Xu and Lu, 2006; Hu *et al.*, 2007; Xie *et al.*, 2007b; Liu *et al.*, 2008; Tang *et al.*, 2008; Yang *et al.*, 2008; Zhou *et al.*, 2008; Yang, 2009). However, it was also misidentified as *E. coelestinum* (Li *et al.*, 2006; Ma *et al.*,

\* Foundation item: Shanghai Landscaping Administrative Bureau (G1024011)

\*\* Xiao-Ling YAN, E-mail: sx\_yxl@163.com, xiaolingyan.zju@gmail.com; Jin-Shuang MA, E-mail: jinshuangma@gmail.com  
Received date: 2010-12-28, Accepted date: 2011-01-10

① Y. L. Chen, T. Kawahara & N. Hind, Asteraceae in Fl. Chin. **20** (unpublished, online: [http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae\\_coauthoring.htm](http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae_coauthoring.htm)).

② 《中国植物志》英文版称其为破坏草, 但国内大多数学者称其为“紫茎泽兰”。

2009; *Conoclinium coelestinum* in *Flora of China*<sup>③</sup>), which is native to North America, and sometimes been classified in the genus *Eupatorium*, especially *Ageratum* (*Ageratina* is the diminutive of *Ageratum*, Schmidt and Schilling, 2000), now has been cultivated and escaped in Guizhou and Yunnan of China.

**2. *Alternanthera bettzickiana***<sup>④</sup> (Regel) G. Nicholson, Ill. Gard. Dict. 59. 1884; B. J. Bao, S. E. Clemants & T. Borsch in Fl. Chin. **5**: 427. 2003. Syn. *Alternanthera versicolor* (Lemaire) Regel, Gartenflora. **18**: 101. 1869 (Amaranthaceae).

Calico plant (锦绣苋), a native species found in South America, is treated as *Alternanthera versicolor* (Shan *et al.*, 2006; Yan *et al.*, 2007). Now it has been cultivated throughout Southeast Asia and China: Anhui, Beijing, Fujian, Gansu, Guangdong, Guangxi, Hainan, Hebei, Heilongjiang, Henan, Hong Kong, Hubei, Hunan, Jiangsu, Jiangxi, Jilin, Inner Mongolia, Liaoning, Macau, Ningxia, Qinghai, Shaanxi, Shandong, Shanghai, Shanxi, Sichuan, Taiwan, Tianjin, Xinjiang, Xizang, Yunnan and Zhejiang.

**3. *Amaranthus blitum*** Linnaeus, Sp. Pl. **1**: 990. 1753; B. J. Bao, S. E. Clemants & T. Borsch in Fl. Chin. **5**: 421. 2003. Syn. *Amaranthus lividus* Linnaeus, Sp. Pl. **1**: 990. 1753 (Amaranthaceae).

Purple amaranth (凹头苋), native to the Mediterranean region and naturalized in other parts of the world including much of eastern North America, is reported as *Amaranthus lividus* (Liu *et al.*, 2006; Xu *et al.*, 2006; Dong and Ye, 2007; Li *et al.*, 2007; Wang *et al.*, 2007a; Zhu *et al.*, 2007; Peng *et al.*, 2008; Song *et al.*, 2009; Niu and Li, 2010; Gao and Cao, 2010) in China: Anhui, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hainan, Hebei,

Heilongjiang, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Jilin, Liaoning, Shaanxi, Shandong, Shanxi, Sichuan, Taiwan, Xinjiang, Yunnan and Zhejiang.

**4. *Amaranthus hybridus*** Linnaeus, Sp. Pl. **2**: 990. 1753; B. J. Bao, S. E. Clemants & T. Borsch in Fl. Chin. **5**: 427. 2003 (Amaranthaceae).

Slim amaranth (绿穗苋) was originally a pioneer plant in eastern North America, and it has been reported in Bhutan, Japan, Laos, Nepal, Sikkim, Vietnam, Europe, North and South America as well, naturalized in many places of warmer climate, and widespread in China: Anhui, Fujian, Guizhou, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Shaanxi, Sichuan and Zhejiang. *Amaranthus chlorostachys* has been suggested as subspecies of *A. hybridus*, while some authors treated it as *A. hybridus* (Chen *et al.*, 2003; Wei *et al.*, 2006a; Yang, 2009).

**5. *Amaranthus tricolor*** Linnaeus, Sp. Pl. **2**: 989. 1753; B. J. Bao, S. E. Clemants & T. Borsch in Fl. Chin. **5**: 420. 2003. Syn. *A. mangostanus* Linnaeus, Cent. Pl. I. **1**: 32. 1755 (Amaranthaceae).

Three-colored amaranth (苋), probably native to tropical Asia, is treated as *Amaranthus mangostanus* (Sun *et al.*, 2009). It is widely cultivated in China: Anhui, Beijing, Chongqing, Fujian, Gansu, Guangdong, Guangxi, Hainan, Hebei, Heilongjiang, Hubei, Hunan, Jiangsu, Jiangxi, Jilin, Inner Mongolia, Liaoning, Macau, Ningxia, Qinghai, Shaanxi, Shandong, Shanghai, Shanxi, Sichuan, Taiwan, Tianjin, Xinjiang, Xizang, Yunnan and Zhejiang.

**6. *Basella alba*** Linnaeus, Sp. Pl. **1**: 272. 1753; D. Q. Lu & M. G. Gilbert in Fl. Chin. **5**: 445. 2003. Syn. *Basella rubra* Linnaeus in Sp. Pl. **1**: 272. 1753 (Basellaceae).

③ Y. L. Chen, T. Kawahara & N. Hind, Asteraceae, in Fl. Chin. 20 (unpublished, online: [http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae\\_coauthoring.htm](http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae_coauthoring.htm)).

④ Flora of North America recorded as *Alternanthera bettzickiana*.

Ceylon spinach (落葵), a pantropical species, is treated as *Basella rubra* (Chen, 2008), widely cultivated in China and naturalized in South China: Gongdong, Guangxi, Hainan, Hong Kong, Macau and Zhejiang.

**7. *Chromolaena odorata*** (Linnaeus) R. M. King & H. Robinson, *Phytologia*. **20**: 204. 1970<sup>⑤</sup>; G. L. Nesom in *Fl. N. Amer.* **21**: 544. 2006. Basionym *Eupatorium odoratum* Linnaeus, *Syst. Nat.*, ed. 10, 1205. 1759 (Asteraceae).

Jack in the bush (飞机草), a species native to North America, from Florida and Texas to Mexico and the Caribbean, widely naturalized in tropical Asia, introduced into Hainan of China during World War II. It is a destructive invasive alien in the field because of very strong ability in propagation and has been naturalized and invasive in many places of China: Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hong Kong, Macau, Sichuan, Taiwan and Yunnan.

*Chromolaena* is here followed *Flora of North America* and *Flora of China*. However, it has frequently been used as *Eupatorium odoratum* in the Chinese works (Li and Xie, 2002; Zeng, 2003; Xu and Qiang, 2004; Wang *et al.*, 2004; Guan *et al.*, 2006; Wei *et al.*, 2006a, b; Xu *et al.*, 2006; Xu and Lu, 2006; An *et al.*, 2007; Ning *et al.*, 2007; Wang *et al.*, 2007b; Xie *et al.*, 2007b; Fan *et al.*, 2008; Liu *et al.*, 2008; Qin *et al.*, 2008; Tang *et al.*, 2008; Wang *et al.*, 2008; Zhao *et al.*, 2008; Zhou *et al.*, 2008; Long, 2009; Ma *et al.*, 2009).

**8. *Cucumis melo*** Linnaeus, *Sp. Pl.* **2**: 1011. 1753<sup>⑥</sup>. Syn. *Cucumis bisexualis* Bull. *Bot. Res.*, Harbin. **4** (2): 126–127, f. 1–6. 1984 (Cucurbitaceae).

Cantaloupe (甜瓜), native to Old World tropics and subtropics, widely cultivated in tropical and

temperate countries, adventive in New World tropics, but wide cultivated and escaped in China: Anhui, Henan, Hunan, Jiangsu, Liaoning, Shandong and Shanxi, but is treated as *Cucumis bisexualis* (Xu *et al.*, 2004; Zhu *et al.*, 2007; He and Ge, 2008). Xu and Qiang (2004) also treated *C. melo* and *C. bisexualis* as two species separately.

**9. *Dysphania ambrosioides*** (Linnaeus) Mosyakin & Clemants, *Ukrayinsk. Bot. Zhurn.* **59**: 382. 2002; G. L. Chu, S. L. Mosyakin & S. E. Clemants in *Fl. Chin.* **5**: 377. 2003. Basionym *Chenopodium ambrosioides* Linnaeus, *Sp. Pl.* **1**: 219. 1753 (Chenopodiaceae).

Mexican-tea (土荆芥), native to tropical America, widely naturalized in tropical, subtropical, and warm-temperate regions of the world, naturalized and often cultivated for medicine in China: Fujian, Guangdong, Guangxi, Hunan, Jiangsu, Jiangxi, Sichuan, Taiwan, Yunnan and Zhejiang. *Dysphania* is accepted here in sensu lato (Mosyakin and Clemants, 2002), including all other “glandular” taxa previously treated in *Chenopodium* subg. The species is widely accepted as *Chenopodium ambrosioides* in China (Li and Xie, 2002; Deng and Lu, 2004; Liu and Qin, 2004; Shi *et al.*, 2004; Wang *et al.*, 2004; Xu and Qiang, 2004; Yan *et al.*, 2004; Hu *et al.*, 2005; Tian *et al.*, 2005; Ding *et al.*, 2006; Guan *et al.*, 2006; Ou and Lu, 2006; Shi *et al.*, 2006; Song *et al.*, 2006; Wei *et al.*, 2006a, b; Wu *et al.*, 2006; Xu and Lu, 2006; Dong and Ye, 2007; Hu *et al.*, 2007; Li *et al.*, 2007; Lin and Wang, 2007; Tan *et al.*, 2007; Wang *et al.*, 2007b; Xie *et al.*, 2007a, b; Zhao, 2007; Zhu *et al.*, 2007; Chen *et al.*, 2008; He and Ge, 2008; Jiang *et al.*, 2008; Li *et al.*, 2008; Liu *et al.*, 2008; Luo, 2008; Peng *et al.*, 2008, 2009; Wang

⑤ Y. L. Chen, T. Kawahara & N. Hind, Asteraceae in *Fl. Chin.* 20 (unpublished, online: [http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae\\_coauthoring.htm](http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae_coauthoring.htm)).

⑥ A. M. Lu & C. Jeffrey, Cucurbitaceae in *Fl. Chin.* 19 (unpublished, online: [http://flora.huh.harvard.edu/china/mss/volume19/Cucurbitaceae-MO\\_reviewing.htm](http://flora.huh.harvard.edu/china/mss/volume19/Cucurbitaceae-MO_reviewing.htm)).

*et al.*, 2008; Yang *et al.*, 2008; Zhou *et al.*, 2008; Ji *et al.*, 2009; Jiang *et al.*, 2009; Li *et al.*, 2009; Ma *et al.*, 2009; Shu *et al.*, 2009; Tang *et al.*, 2008; Sun *et al.*, 2009; Xu *et al.*, 2009; Yang and Chen, 2009; Yu *et al.*, 2009; Zeng *et al.*, 2009; Zhang and Zhang, 2009; Ge *et al.*, 2010; He *et al.*, 2010; Qi *et al.*, 2010; Wang, 2010; Zhang *et al.*, 2010).

**10. *Euphorbia maculata*** Linnaeus, Sp. Pl. **1**: 455. 1753; J. S. Ma & M. G. Gilbert in Fl. Chin. **11**: 296. 1997. Syn. *Euphorbia supina* Rafinesque in Amer. Monthly Mag. & Crit. Rev. **2** (2): 119. 1817 (Euphorbiaceae).

Spotted sandmat (斑地锦), a species native to North America and naturalized in China: Hebei, Henan, Hubei, Jiangsu, Jiangxi, Taiwan and Zhejiang, is treated as *Euphorbia supina* (Liu *et al.*, 2002; Li *et al.*, 2008; Yang *et al.*, 2009; Dong *et al.*, 2010).

**11. *Ipomoea alba*** Linnaeus, Sp. Pl. **1**: 161. 1753; R. C. Fang & G. Staples in Fl. Chin. **16**: 309. 1995. Syn. *Calonyction aculeatum* (Linnaeus) House, Bull. Torrey Bot. Club. **31** (11): 590. 1904 (Convolvulaceae).

Tropical white morning-glory (月光花), a species of night-blooming morning-glory, native to tropical and subtropical regions of the New World, from northern Argentina north to Mexico and Florida, cultivated and also naturalized in Indonesia, Japan (Ryukyu Islands), Malaysia, Myanmar, Nepal, New Guinea, Philippines, Sri Lanka, Thailand and Pacific Islands, was used as *Calonyction aculeatum* (Yan *et al.*, 2007; Chen, 2008; Zeng *et al.*, 2009; Qi *et al.*, 2010). This species is widely cultivated as an ornamental plant for its flowers in China: Guangdong, Guangxi, Hainan, Jiangsu, Jiangxi, Shaanxi, Sichuan, Yunnan and Zhejiang.

**12. *Ipomoea nil*** (Linnaeus) Roth, Catal. Bot. **1**: 36. 1797. R. C. Fang & G. Staples in Fl. Chin.

**16**: 305. 1995. Basionym *Convolvulus nil* Linnaeus, Sp. Pl. ed. 2. **1**: 219. 1762. Syn. *Ipomoea hederacea* (Linnaeus) Jacquin, C. B. Clarke in Hook. f. Fl. Brit. Ind. **4**: 199. 1883; *Pharbitis nil* (Linnaeus) Choisy in Mém. Soc. Phys. Genève. **6**: 439–440. 1833 (Convolvulaceae).

Whiteedge morning-glory (牵牛) has been treated either as *Pharbitis hederacea* (Ma *et al.*, 2009; Qi *et al.*, 2010) or *Ipomoea hederacea* (Chen *et al.*, 2006), or was frequently recorded as *Pharbitis nil* (Yan *et al.*, 2004, 2005; Chen, 2005; Ding *et al.*, 2006; Qu *et al.*, 2006a, b; Jiang *et al.*, 2008; Zhou *et al.*, 2008; Chen, 2008; Zeng *et al.*, 2009). Native to South America, it now nearly cultivated as an attractive ornamental plant in many pantropical places, and the descendents of garden escapes now grows in China: Anhui, Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hebei, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Inner Mongolia, Ningxia, Shaanxi, Shandong, Shanxi, Sichuan, Taiwan, Xizang, Yunnan and Zhejiang.

**13. *Ipomoea quamoclit*** Linnaeus, Sp. Pl. **1**: 159–160. 1753; R. C. Fang & G. Staples in Fl. Chin. **16**: 301. 1995. Syn. *Quamoclit pennata* (Desr.) Bojer, Hortus Maurit. 224. 1837 (Convolvulaceae).

Cypressvine (茛萝), native to tropical regions of the New World from northern South America to Mexico, is reported as *Quamoclit pennata* (Li, 1992; Qu *et al.*, 2006a, b; Wang, 2008; Qi *et al.*, 2010). It is widely cultivated as ornamental plant throughout the tropics and also outside of tropical regions, cultivated and escapes from gardens in China: Guangdong, Hebei, Liaoning, Shaanxi and Zhejiang.

**14. *Lablab purpureus*** (Linnaeus) Sweet, Hort. Brit. 481. 1826; T. L. Wu & M. Thulin in Fl. Chin. **10**: 253. 2010. Syn. *Dolichos lablab* Linnaeus in Fl. Galápagos Isl. i-xx, 1–998. 1971 (Fabaceae).

Hyacinth Bean (扁豆), native to Africa, cultivated throughout the tropics, widely cultivated in China: Beijing, Chongqing, Hainan, Hebei, Hong

Kong, Shaanxi and Zhejiang, is treated as *Dolichos lablab* (Liu *et al.*, 2002; Sun *et al.*, 2009).

**15. *Leucaena leucocephala*** (Lamarck) de Wit, Taxon **10**: 54. 1961; T. L. Wu & I. C. Nielsen in Fl. Chin. **10**: 53. 2010. Basionym *Mimosa leucocephala* Lamarck, Encycl. **1**: 12. 1783. Syn. *Leucaena glauca* Benth, J. Bot. (Hooker). **4** (32): 416. 1842 (Fabaceae).

White leadtree (银合欢) originally from tropical America, widely distributed in tropical and subtropical regions, cultivated and naturalized for use as green manure and fodder in China: Fujian, Guangdong, Guangxi, Guizhou, Hainan, Taiwan and Yunnan, is treated as *Leucaena glauca* (Peng *et al.*, 2008; Qin *et al.*, 2008).

**16. *Medicago polymorpha*** Linnaeus, Sp. Pl. **2**: 779. 1753; Z. Wei & M. A. Vincent in Fl. Chin. **10**: 557. 2010. Syn. *Medicago hispida* in Fruct. Sem. Pl. **2**: 349, pl. 155. 1791 (Fabaceae).

Burclover (南苜蓿), annual or biennial herbs, native at least to N Africa, SW Asia, and S Europe, widely cultivated or escaped to elsewhere fields, including China: Anhui, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hainan, Hubei, Hunan, Jiangsu, Jiangxi, Shaanxi, Sichuan, Taiwan, Yunnan and Zhejiang. However, it was commonly accepted as *Medicago hispida* in China (Liu and Qin, 2004; Xu and Qiang, 2004; Lu *et al.*, 2005; Yi *et al.*, 2005; Shi *et al.*, 2006; Wang *et al.*, 2007a; Chen *et al.*, 2008; He and Ge, 2008; Wang, 2008; Zhou *et al.*, 2008; Ji *et al.*, 2009; Ma *et al.*, 2009; Ge *et al.*, 2010; Wang, 2010).

**17. *Melinis repens*** (Willdenow) Zizka, Biblioth. Bot. **138**: 55. 1988; S. L. Chen & S. M. Phillips in Fl. Chin. **22**: 539, 2006. Basionym *Saccharum repens* Willdenow, Sp. Pl., ed. 4, **1**: 322. 1797. Syn. *Rhy-*

*nchelytrum repens* (Willdenow) C. E. Hubbard, Bull. Misc. Inform. Kew **1934** (3): 110. 1934 (Poaceae).

Rose Natal grass (红毛草), native to Africa, naturalized in China: Fujian, Guangdong, Hainan, Hong Kong, Macau, Taiwan and Yunnan, is a polymorphic and pantropical weed, recognizable by its pink, fluffy panicles. However, it was widely accepted as *Rhynchelytrum repens* in China (Li and Xie, 2002; Wang *et al.*, 2004; Chen, 2005; Yan *et al.*, 2004, 2005; Li *et al.*, 2006; An *et al.*, 2007; Fan *et al.*, 2008; Jiang *et al.*, 2008; Qin *et al.*, 2008; Lin *et al.*, 2008; Wang *et al.*, 2008; Yang and Chen, 2009).

**18. *Mimosa bimucronata*** (Candolle) O. Kuntze, Revis. Gen. Pl. **1**: 198. 1891; T. L. Wu & I. C. Nielsen in Fl. Chin. **10**: 53, 2010. Basionym *Aca-cia bimucronata* Candolle, Prodr. **2**: 469. 1825. Syn. *Mimosa sepiaria* Benth. Britton, N. Amer. Fl. **23** (3): 137–194. 1928 (Fabaceae).

Smooth pod mimic (光荚含羞草), native to South America (NE Argentina and E Brazil), introduced to Guangdong, escaped in spare forests of Guangxi, Hainan and Yunnan of China. It has been treated as *Mimosa sepiaria* (Ding *et al.*, 2006; Xie *et al.*, 2007b; Ye *et al.*, 2008). Fan *et al.* (2008) also used as “*Mimosa sphyria*”, the name first used by Liu (2006) in his doctoral dissertation, but never existed in nomenclature.

**19. *Mukia maderaspatana*** (Linnaeus) M. J. Roemer, Fam. Nat. Syn. Monogr. **2**: 47. 1846<sup>⑦</sup>. Syn. *Coccinia cordifolia* (Linnaeus) Cogniaux, Monogr. Phan. **3**: 529. 1881 (Cucurbitaceae).

Ivy gourd (帽儿瓜), native to old tropical and subtropical and Australia, is accepted as *Coccinia cordifolia* in China: Guangdong, Guangxi, Guizhou, Taiwan and Yunnan (Xu and Qiang, 2004; Wang *et al.*, 2007b).

⑦ A. M. Lu & C. Jeffrey, Cucurbitaceae in Fl. Chin. 19 (unpublished, online; [http://flora.huh.harvard.edu/china/mss/volume19/Cucurbitaceae-MO\\_reviewing.htm](http://flora.huh.harvard.edu/china/mss/volume19/Cucurbitaceae-MO_reviewing.htm)).

**20. *Oenothera stricta*** Ledebour ex Link, Enum. Pl. Hort. Berol. **1**: 377. 1821; J. R. Chen, P. C. Hoch & W. L. Wagner in Fl. Chin. **13**: 426. 2007 (Onagraceae).

Chilean evening primrose (待宵草), native to South America (Chile and Argentina), naturalized in Africa, SW Asia, Australia, Europe, North America and Pacific islands, usually escaped from cultivation and sometimes cultivated for its relatively large, attractive flowers, often becomes naturalized in China and elsewhere: Fujian, Guangxi, Guizhou, Hubei, Jiangxi, Shaanxi, Shandong, Sichuan, Taiwan and Yunnan.

Many reports from China have been determined as *Oenothera odorata* Jacquin (Yi *et al.*, 2005; Zhao and Dong, 2007; Chen, 2008; Qi *et al.*, 2010), however, the later, also native to S South America but rarely, if ever, naturalized elsewhere, differs from *O. stricta* by having narrower leaves, larger petals that lacks a red spot at the base, shorter floral tubes, and bracts mostly longer than the capsules they subtend.

**21. *Opuntia dillenii*** (Ker Gawler) Haworth, Suppl. Pl. Succ. 79. 1819; Z. Y. Li & N. P. Taylor in Fl. Chin. **13**: 210. 2007. Basionym *Cactus dillenii* Ker Gawler, Bot. Reg. **3**: t. 255. 1818. Syn. *Opuntia stricta* (Haworth) Haworth var. *dillenii* (Ker Gawler) L. Benson, in Cact. Succ. J. (Los Angeles), **41** (3): 126. 1969 (Cactaceae).

Erect prickly pear (仙人掌), native to the Caribbean region, widely introduced and naturalized in tropical regions, is treated as *Opuntia stricta* var. *dillenii* in China (Xu *et al.*, 2006; An *et al.*, 2007; Xie *et al.*, 2007b; Fan *et al.*, 2008; Qin *et al.*, 2008; Tang *et al.*, 2008; Wang *et al.*, 2008; Yang *et al.*, 2008; Yan and Chen, 2009). The species was first recorded in China in 1702, and now widely spread in China: Chongqing, Fujian,

Guangdong, Guangxi, Hainan, Hong Kong, Hunan, Macau, Shandong, Yunnan and Zhejiang.

**22. *Paspalum distichum*** Linnaeus, Syst. Nat., ed. 10, **2**: 855. 1759; S. L. Chen & S. M. Phillips in Fl. Chin. **22**: 528. 2006. Syn. *Paspalum paspalodes* (Michaux) Scribner, Mem. Torrey Bot. Club **5** (3): 29. 1894 (Poaceae).

Knot grass (双穗雀稗), a monocot, is native to California and is also found elsewhere in North America, tropical and warm-temperate regions of the world, including China: Anhui, Fujian, Guangxi, Guizhou, Hainan, Henan, Hong Kong, Hubei, Hunan, Jiangsu, Shandong, Sichuan, Taiwan, Yunnan and Zhejiang. The length of the lower glumes is very variable in this species, even within a single raceme. It is reported as *Paspalum paspaloides* (Guan *et al.*, 2006).

**23. *Praxelis clematidea*** R. M. King & H. Robinson, Phytologia. **20**: 194. 1970<sup>⑧</sup>. Syn. *Eupatorium catarium* Veldkamp, Gard. Bull. Singapore **51**: 121. 1999 (Asteraceae).

Praxelis (假臭草), widespread in South America, becoming weedy in many cultivated areas, has been introduced and becoming naturalized in Northern Australia and Asia (Fujian, Guangdong, Guangxi, Hainan, Hong Kong, Macau and Taiwan of China). It was commonly accepted as *Eupatorium catarium* (Li and Xie, 2002; Wang *et al.*, 2004; Yan *et al.*, 2004, 2005; Chen, 2005; Li *et al.*, 2006; Ou and Lu, 2006; Zheng *et al.*, 2006; An *et al.*, 2007; Cao *et al.*, 2007; Ning *et al.*, 2007; Wang *et al.*, 2007b; Chen *et al.*, 2008; Fan *et al.*, 2008; Jiang *et al.*, 2008; Lin *et al.*, 2008; Qin *et al.*, 2008; Wang *et al.*, 2008; Yang and Chen, 2009; Zeng *et al.*, 2009).

**24. *Senna tora*** (Linnaeus) Roxburgh, Fl. Ind., ed. **2**: 340. 1832; T. C. Chen, D. X. Zhang & K.

⑧ Y. L. Chen, T. Kawahara & N. Hind, Asteraceae in Fl. Chin. 20 (unpublished, online: [http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae\\_coauthoring.htm](http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-AGH-Eupatorieae_coauthoring.htm)).



Larsen in Fl. Chin. **10**: 32. 2010. Basionym *Cassia tora* Linnaeus, Sp. Pl. **1**: 376. 1753. Syn. *C. obtusifolia* L., Ann. Missouri Bot. Gard. **38** (1): 1-94. 1951 (Fabaceae).

Sickle senna (决明) has been widely accepted either as *Cassia tora* (Ng and Richard, 2002; Deng and Lu, 2004; Tian and Lü, 2004; Wang *et al.*, 2004; Xu and Qiang, 2004; Yan *et al.*, 2004; Ding *et al.*, 2006; Guan *et al.*, 2006; Liu *et al.*, 2006; Qi and Xu, 2006; Shi *et al.*, 2006; Song *et al.*, 2006; Wei *et al.*, 2006a; An *et al.*, 2007; Qin *et al.*, 2007; Wang *et al.*, 2007; Chen *et al.*, 2008; Fan *et al.*, 2008; He and Ge, 2008; Jiang *et al.*, 2008; Luo and Liu, 2008; Peng *et al.*, 2008, 2009; Qin *et al.*, 2008; Wang, 2008; Zhou *et al.*, 2008; Chen *et al.*, 2009; Li *et al.*, 2009; Ma *et al.*, 2009; Shu *et al.*, 2009; Xu *et al.*, 2009; Zeng *et al.*, 2009; Zhang and Zhang, 2009; Zhang *et al.*, 2010), or *Cassia obtusifolia* (He *et al.*, 2010). It is native to tropical America and widely cultivated in the tropics, subtropics and China: Anhui, Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hebei, Hong Kong, Hunan, Jiangsu, Liaoning, Macau, Shandong, Shanghai, Sichuan, Taiwan, Yunnan and Zhejiang.

**25. *Solanum aculeatissimum*** Jacquin, Collectanea **1**: 100. 1787; Z. Y. Zhang, A. M. Lu & W. G. D'Arcy in Fl. Chin. **17**: 323. 1994. Syn. *Solanum khasianum* C. B. Clarke, Fl. Brit. India **4** (10): 234. 1883 (Solanaceae).

Indian nightshade (喀西茄), probably native to Brazil, widespread in old tropical and China: Fujian (Xiamen), Guangxi, Guizhou, Hunan, Jiangxi, Sichuan, Xizang, Yunnan and Zhejiang (Pingyang), is recorded as *Solanum khasianum* (Ding *et al.*, 2006; Qin *et al.*, 2008).

**26. *Solidago canadensis*** Linnaeus, Sp. Pl. **2**: 879.

1753<sup>⑨</sup>. Syn. *Solidago altissima* Linnaeus, Monogr. Syst. Bot. Missouri Bot. Gard. **85**: i-xlii, 1-2666. 2001 (Asteraceae).

Canada goldenrod (加拿大一枝黄花), an ornamental and decorative plant, native to North America, widely cultivated in gardens and parks in China: Anhui, Beijing, Fujian, Gansu, Guangxi, Hebei, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Jilin, Liaoning, Shaanxi, Shanghai, Shanxi, Sichuan, Yunnan and Zhejiang. It is one of the worst weeds and most widespread invasive species in China, but is treated as *Solidago altissima* (Peng *et al.*, 2008).

**27. *Spermacoce latifolia*** Aublet, Hist. Pl. Guiane **1**: 55, t. 19/1. 1775. Syn. *Borreria latifolia* (Aublet) K. Schum in Fl. Bras. **6** (6): 61, t. 80. 1888 (Rubiaceae).

Button weed (阔叶丰花草), a native species found in South America, introduced to Guangdong, naturalized in wastelands of China: Fujian, Guangdong, Guangxi, Hainan, Hong Kong, Hunan, Macau, Taiwan, Yunnan and Zhejiang, is treated as *Borreria latifolia* (Li *et al.*, 2006; Zheng *et al.*, 2006; An *et al.*, 2007; Wang *et al.*, 2008).

**28. *Symphytum officinale*** Linnaeus, Sp. Pl. **1**: 136. 1753; G. L. Zhu, H. Riedl & R. Kamelin in Fl. Chin. **16**: 359. 1995. Syn. *Symphytum peregrinum* Ledeb., Ind. Sem. Hort. Dorp. **4**. 1820 (Boraginaceae).

Comfrey (聚合草), native to Europe, is known elsewhere, including North America, was introduced in 1963 in China: Fujian, Hebei, Liaoning, Taiwan and Xinjiang as green forage for livestock, but was reported as *Symphytum peregrinum* (Li, 1981, 1992; Qu *et al.*, 2006a, b).

**29. *Veronica polita*** Fries, Novit. Fl. Suec. **5**: 63. 1817. D. Y. Hong & M. A. Fischer in Fl. Chin.

⑨ Y. L. Chen, L. Brouillet & J. C. Semple, Asteraceae in Fl. Chin. 20 (unpublished online; [http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-MO-Astereae\\_coauthoring.htm](http://flora.huh.harvard.edu/china/mss/volume20/Asteraceae-MO-Astereae_coauthoring.htm)).

**18: 70.** 1998. Syn. *Veronica didyma* Tenore., So-sud. Rast. SSSR. 509. 1981 (Scrophulariaceae).

Grey field-speedwell (婆婆纳), native in SW Asia and naturalized over most of the world, is treated as *Veronica didyma* (Liu and Qin, 2004; Tian *et al.*, 2005; Xie *et al.*, 2007b; Zhu *et al.*, 2007; Li *et al.*, 2008; Shu *et al.*, 2009; Zhang, 2009), widely spread in China: Anhui, Beijing, Fujian, Gansu, Guizhou, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Qinghai, Shaanxi, Sichuan, Taiwan, Xinjiang, Yunnan and Zhejiang.

**30. *Waltheria indica*** Linnaeus, Sp. Pl. **2**: 673. 1753; Y. Tang, M. G. Gilbert & L. J. Don in Fl. Chin. **12**: 321. 2007. Syn. *Waltheria americana* Linnaeus in Sp. Pl. **2**: 673. 1753 (Sterculiaceae).

Uhaloa (蛇婆子), pantropical species, is believed to have originated from the Neotropics, and is treated as *Waltheria americana* (An *et al.*, 2007; Wang *et al.*, 2008) in south China: Fujian, Guangdong, Guangxi, Hainan, Hong Kong, Macau, Taiwan and Yunnan.

## Reference:

- An F (安锋), Kan LY (阚丽艳), Xie GS (谢贵水) *et al.*, 2007. Alien invasion plants in Hainan island and contral countermeasures [J]. *Journal of Northwest Forestry University* (西北林学院学报), **22** (5): 198—206
- Chen HB (陈恒彬), 2005. The harmful exotic plants in Xiamen area [J]. *Subtropical Plant Science* (亚热带植物科学), **34** (1): 50—55
- Chen MS (陈模舜), 2008. Studies on flora composition and distribution characteristics of alien plants in Taizhou [J]. *Journal of Taizhou University* (台州学院学报), **30** (6): 43—48
- Chen QX (陈秋霞), Wei CQ (韦春强), Tang SC (唐赛春) *et al.*, 2008. The alien invasive plants in Guilin city, Guangxi Zhuangzu Autonomous region [J]. *Subtropical Plant Science* (亚热带植物科学), **37** (3): 55—58
- Deng XC (邓晰朝), Lu X (卢旭), 2004. A preliminary study on the invasive plants in Yizhou city [J]. *Journal of Hechi Normal College* (河池师专学报), **24** (2): 72—74
- Ding L (丁莉), Du F (杜凡), Zhang DC (张大才), 2006. A study of alien invasive plant in Yunnan Province [J]. *Journal of West China Forestry Science* (西部林业科学), **35** (4): 98—103, 108
- Dong DP (董东平), Ye YZ (叶永忠), 2007. Studies on flora alien invasive plants and disastrous mechanisms in Henan [J]. *Henan Science* (河南科学), **25** (5): 765—769
- Dong HY (董红云), Li Y (李亚), Wang Q (汪庆) *et al.*, 2010. Investigation and analysis on alien invasive plants in three Nature Reserves of Jiangsu Province [J]. *Journal of Plant Resources and Environment* (植物资源与环境学报), **19** (1): 86—91
- Fan ZW (范志伟), Shen YD (沈奕德), Liu LZ (刘丽珍), 2008. List of invasive alien weeds in Hainan, China [J]. *Chinese Journal of Tropical Crops* (热带作物学报), **29** (6): 781—792
- Fang RC, Staples G, 1995. Convolvulaceae [A]. In *Flora of China* [M]. Beijing: Science Press, St. Louis: Missouri Botanical Garden Press, **16**: 271—327
- Gao Y (高燕), Cao W (曹伟), 2010. Situation and countermeasure of alien invasive plants in Northeast China [J]. *Journal of the Graduate School of the Chinese Academy of Sciences* (中国科学院研究生院学报), **27** (2): 191—198
- Ge G (葛刚), Li EX (李恩香), Wu HP (吴和平) *et al.*, 2010. Invasive plants in the National Nature Reserve of Lake Poyang [J]. *Journal of Lak. Sciences* (湖泊科学), **22** (1): 93—97
- Guan ZB (管志斌), Deng WH (邓文华), Huang ZL (黄志玲) *et al.*, 2006. A preliminary investigation on the alien invasive plants in Xishuangbanna [J]. *Tropical Agricultural Science & Technology* (热带农业科技), **29** (4): 35—38
- He DM (何冬梅), Lu XZ (鲁小珍), Yi XG (伊贤贵) *et al.*, 2010. Investigation and prevention measures on the invasive alien plants in Bengbu city of Anhui Province [J]. *Journal of Anhui Agricultural Science* (安徽农业科学), **38** (6): 3081—3083
- He JQ (何家庆), Ge JL (葛结林), 2008. The present situation of alien invasive plants in Anhui Province and a comparison with the other provinces in China [J]. *Journal of Anhui University* (Natural Science Edition) (安徽大学学报 (自然科学版)), **32** (2): 82—89
- Hu G (胡刚), Zhang ZH (张忠华), Dong JT (董金廷) *et al.*, 2005. A Preliminary study on the alien invasive plants in Huai-bei region of Anhui province [J]. *Journal of Hefei University* (Natural Sciences) (合肥学院学报 (自然科学版)), **15** (2): 41—45
- Hu TY (胡天印), Jiang HW (蒋华伟), Fang F (方芳) *et al.*, 2007. Preliminary survey on exotic invasive plants in Lushan Mountain scenic spot [J]. *Jiangxi Forestry Science and Technology* (江西林业科技), **3**: 19—23
- Huang GQ (黄国勤), Huang QP (黄秋萍), 2006. The status Quo, Harm and countermeasure of biology invasive in Jiangxi Province [J]. *Meteorology and Disaster Reduction Research* (气象与减灾研究), **29** (1): 51—55
- Ji CF (季春峰), Wang Z (王智), Qian P (钱萍), 2009. Primary research on invasive plants in Jiangxi Province [J]. *Hubei Forestry Science and Technology* (湖北林业科技), **3**: 27—31



- Jiang GC (姜罡丞), Wang K (王珂), Dong DP (董东平), 2009. Research and risk assessment of alien invasive plants of Xuchang suburb [J]. *Journal of Henan Normal University (Natural Science)* (河南师范大学学报(自然科学版)), **37** (6): 168—170
- Jiang QC (蒋谦才), Lin ZM (林正眉), Li L (李荔) *et al.*, 2008. An investigation on the invasive plants in Zhongshan, Guangdong Province [J]. *Guangdong Forestry Science and Technology* (广东林业科技), **24** (2): 54—58
- Li HS (李海生), Zhong HL (钟化龙), Liu G (刘广) *et al.*, 2006. A preliminary study on the alien invasive plant species in Baiyun Mountain, Guangzhou [J]. *Journal of Guangdong Education Institute* (广东教育学院学报), **28** (3): 65—68
- Li HX (李惠欣), 2008. A preliminary study on invasive plants of Hebei Hengshui Lake Natural Reserve [J]. *Wetland Science & Management* (湿地科学与管理), **4** (2): 51—53
- Li SC (李顺才), Liu LX (刘连新), Hu XD (胡晓东) *et al.*, 2009. Investigation and analysis of alien invasions in Qinhuangdao [J]. *Journal of Hebei Normal University of Science & Technology* (河北科技师范学院学报), **23** (3): 13—17
- Li SX (李书心), 1981. *Symphytum* [A]. In Liou TN ed, *Flora Plantarum Herbacearum Chinae Boreali-Orientalis* [M]. Beijing: Science Press, **7**: 150
- Li SX (李书心), 1992. *Symphytum* [A]. *Flora Liaoningica* [M]. Shenyang: Liaoning Science and Technology Press, **2**: 167
- Li XW (李乡旺), Hu ZH (胡志浩), Hu XL (胡晓立) *et al.*, 2007. A preliminary study on the exotic invasive plants in Yunnan Province [J]. *Journal of Southwest Forestry College* (西南林学院学报), **27** (6): 5—10
- Li Y (李亚), Yao G (姚淦), Deng F (邓飞) *et al.*, 2008. Preliminary investigation and analysis on exotic spermatophyte in Jiangsu Province [J]. *Journal of Plant Resources and Environment* (植物资源与环境学报), **17** (4): 55—60
- Li ZY (李振宇), Xie Y (解焱), 2002. Invasive Alien Species in China [M]. Beijing: China Forestry Publishing House, 99—189
- Lin HH (林鸿辉), Pan YH (潘永华), Dai SP (代色平) *et al.*, 2008. Analysis on resources of garden plants in the parks of Macau [J]. *Landscape Science & Technology* (园林科技), **30** (4): 5—8
- Liu F (刘峰), Tao GD (陶国达), Wang DS (王东升), 2008. Investigation on the alien invasive plants in Nabanhe Nature Reserve and precautionary measures [J]. *Forest Inventory and Planning* (林业调查规划), **33** (6): 112—117
- Liu GD (刘国道), 2006. Feed Plants Resources in Hainan and Studies on Substitute Coloring Agent from Plants Used for Poultry [M]. Hainan: Danzhou South China University of Tropical Agriculture
- Liu QN (刘庆年), Liu JZ (刘俊展), Liu JT (刘京涛) *et al.*, 2006. Preliminary study on the alien invasive and harmful biotices in the Yellow River delta [J]. *Journal of Shandong Agricultural University (Natural Science)* (山东农业大学学报(自然科学版)), **37** (4): 581—585
- Liu QR (刘全儒), Yu M (于明), Zhou YL (周云龙), 2002. A preliminary study on the invasive plants in Beijing [J]. *Journal of Beijing Normal University (Natural Science)* (北京师范大学学报(自然科学版)), **38** (3): 399—404
- Liu SX (刘胜祥), Qin W (秦伟), 2004. Study on the current status of invasive plants in Hubei Province [J]. *Journal of Central China Normal University (Natural Science)* (华中师范大学学报(自然科学版)), **38** (2): 223—227
- Lin Y (林杨), Wang DM (王德明), 2007. Study on the important value of exotic plants community in Changsha of Hunan Province [J]. *Journal of Anhui Agricultural Science* (安徽农业科学), **35** (4): 1102—1103
- Long YB (龙永彬), 2009. Distribution of five species of alien invasive plants in Guangdong Province [J]. *Journal of Hebei Agricultural Sciences* (河北农业科学), **13** (5): 84—85
- Lu SF (卢少飞), Liu SX (刘胜祥), Fang YP (方元平), 2005. A preliminary research of invasive species around Xingdoushan National Nature Reserve, Huibei [J]. *Journal of Huanggang Normal University* (黄冈师范学院学报), **25** (3): 48—52
- Luo MY (罗明永), 2008. Preliminary study on main foreign invasion species in Fujian [J]. *Journal of Fujian Forestry Science and Technology* (福建林业科技), **35** (2): 167—170
- Luo Y (罗艳), Liu AH (刘爱华), 2008. Preliminary research on alien invasive plants of Qingdao [J]. *Shandong Science* (山东科学), **21** (4): 19—23
- Ma DW (马丹炜), Yu SH (于树华), Mao RB (毛汝兵), 2009. Primary study of the areal types of field vascular weeds in Sichuan Province [J]. *Journal of Sichuan Normal University (Natural Science)* (四川师范大学学报(自然科学版)), **32** (1): 93—97
- Mosyakin SL, Clemants SE, 2002. New nomenclatural combinations in *Dysphania* R. Br. (Chenopodiaceae): Taxa occurring in North America [J]. *Ukrayins'k Bot Zhurn*, n. s. **59**: 380—385
- Ng SC (吴世捷), Richard C (高力行), 2002. The bad biodiversity: alien plant species in Hong Kong [J]. *Biodiversity Science* (生物多样性), **10** (1): 109—118
- Ning ZY (宁昭玉), Hu SQ (胡树泉), Wei YZ (魏远竹) *et al.*, 2007. Present situation of alien species invasion and its effects on economy, society and ecology in Fujian Province [J]. *Entomological Journal of East China* (华东昆虫学报), **16** (4): 304—309
- Niu YL (牛玉璐), Li JM (李建明), 2010. Investigation and study on invasive plants in farmland of Hengshui [J]. *Hubei Agricultural Sciences* (湖北农业科学), **49** (4): 881—883
- Ou J (欧健), Lu CY (卢昌义), 2006. The research of alien plants risk assessment system in Xiamen municipality [J]. *Journal of Xiamen University (Natural Science)* (厦门大学学报(自然科学版)), **45** (6): 883—888
- Peng YL (彭友林), Wang Y (王云), Zhou GQ (周国庆) *et al.*, 2008. Study on the species, distribution and damage of the inva-

- sive alien harmful plants in Dongting area [J]. *Journal of Anhui Agricultural Science* (安徽农业科学), **36** (3): 1114—1116
- Peng YL (彭友林), Wang ZH (王朝晖), Wang Y (王云) *et al.*, 2009. Studies on species, distribution and hazards of exotic harmful plants in Changde city [J]. *Hubei Agricultural Sciences* (湖北农业科学), **48** (8): 1906—1909
- Qi SY (齐淑艳), Xu WD (徐文铎), 2006. Study on types composition and distribution characteristics of alien invasive plants in Liaoning [J]. *Journal of Liaoning Forestry Science & Technology* (辽宁林业科技), **3**: 11—15
- Qi YZ (祁云枝), Du YJ (杜勇军), Zhang Y (张莹), 2010. A investigation of alien invasive plants in Xi'an area [J]. *Chinese Agricultural Science Bulletin* (中国农学通报), **26** (5): 223—227
- Qin WH (秦卫华), Wang Z (王智), Xu WG (徐网谷) *et al.*, 2008. Investigation and analysis on alien invasive plants in three National Nature Reserves in Hainan Province [J]. *Journal of Plant Resources and Environment* (植物资源与环境学报), **17** (2): 44—49
- Qu B (曲波), Lü GZ (吕国忠), Yang H (杨红) *et al.*, 2006a. Report about alien invasive plants in Liaoning Province [J]. *Journal of Shenyang Agricultural University* (沈阳农业大学学报), **37** (4): 587—592
- Qu B (曲波), Lü GZ (吕国忠), Yang H (杨红) *et al.*, 2006b. Preliminary report about alien invasive plants in Liaoning Province [J]. *Liaoning Agricultural Sciences* (辽宁农业科学), **4**: 22—25
- Shan JL (单家林), Yang FC (杨逢春), Zheng XQ (郑学勤), 2006. Exotic plants in Hainan Province [J]. *Subtropical Plant Science* (亚热带植物科学), **35** (3): 39—44
- Shi LC (石亮成), Shi G (石钢), Yi QL (易巧玲) *et al.*, 2009. The countermeasure research for investigation and eradication of alien invasive plant in Liuzhou [J]. *Journal of Guangxi Academy of Sciences* (广西科学院学报), **25** (3): 178—182
- Shi SZ (石胜璋), Tian MJ (田茂洁), Liu YC (刘玉成), 2004. Investigation and study on the alien invasive plants in Chongqing [J]. *Journal of Southwest China Normal University* (Natural Science Edition) (西南师范大学学报 (自然科学版)), **29** (5): 863—866
- Shi Y (石瑛), Xie SL (谢树莲), Wang HL (王惠玲), 2006. Study on the exotic invasive plants in Shanxi Province [J]. *Journal of Tianjin Normal University* (Natural Science Edition) (天津师范大学学报 (自然科学版)), **26** (4): 23—27
- Shu MY (舒美英), Cai JG (蔡建国), Fang BS (方宝生), 2009. Situation and control measures of alien invasive plants in Xixi national wetland park, Hangzhou city, Zhejiang Province [J]. *Journal of Zhejiang Forestry College* (浙江林学院学报), **26** (5): 755—761
- Song N (宋楠), Song YJ (宋亚团), Wang RQ (王仁卿) *et al.*, 2006. Analysis of the present situation and control strategies of the agricultural alien plants in Shandong Province [J]. *Shandong Science* (山东科学), **19** (3): 15—21
- Song XL (宋小玲), Cao F (曹飞), He YH (何云核) *et al.*, 2009. A survey of invasive alien plant species in Dinghushan National Nature Reserve [J]. *Journal of Zhejiang Forestry College* (浙江林学院学报), **26** (4): 538—543
- Sun J (孙娟), Yang GF (杨国锋), Chen YC (陈玉成) *et al.*, 2009. Exotic plants in the Jinpo Mountain Nature Reserve: species ascertain and their distribution [J]. *Acta Prataculturae Sinica* (草业学报), **18** (3): 34—42
- Tang SC (唐赛春), Lü SH (吕仕洪), He CX (何成新) *et al.*, 2008. The alien invasive plants in Guangxi [J]. *Guihaia* (广西植物), **28** (6): 775—779
- Tian CY (田朝阳), Li JZ (李景照), Xu JW (徐景文) *et al.*, 2005. Studies on the alien invasive plants and their control in Henan Province [J]. *Journal of Henan Agricultural Sciences* (河南农业科学), **1**: 31—34
- Tian JY (田家怡), Lü CX (吕传笑), 2004. Harmful AIS in Shandong and their geographic distribution [J]. *Journal of Binzhou Teachers College* (滨州师专学报), **20** (4): 42—46
- Tan MH (谭密红), Deng HP (邓洪平), Tang L (唐兰) *et al.*, 2007. Investigation and study on the alien invasive plants in Xishui of Chongqing [J]. *Journal of Chongqing Three Gorges University* (重庆三峡学院学报), **23** (3): 16—19
- Wang FG (王发国), Xing FW (邢福武), Ye HG (叶华谷) *et al.*, 2004. Preliminary study on invasive species in Macau [J]. *Acta Scientiarum Naturalium Universitatis (Sunyatseni)* (中山大学学报 (自然科学版)), **43** (Suppl.): 106—110
- Wang N (王宁), 2010. Study on the invasiveness and clonality of invasive alien plant in Jiangxi Province [J]. *Journal of Jinggangshan University* (Natural Science) (井冈山大学学报 (自然科学版)), **31** (2): 108—112
- Wang NX (王嫩仙), 2008. Simple study on plants of adventitious invasion in Hangzhou city [J]. *Forest Inventory and Planning* (林业调查规划), **33** (3): 125—128
- Wang W (王伟), Zhang XM (张先敏), Sha LH (沙林华) *et al.*, 2007a. Roster of alien invasive perilous species of animals and plants in Hainan island (I) [J]. *Chinese Journal of Tropical Agriculture* (热带农业科学), **27** (4): 58—64
- Wang XF (汪小飞), Cheng YH (程轶宏), Zhao CH (赵昌恒) *et al.*, 2007b. Studies on alien plants in Huangshan city [J]. *Journal of Jiangsu Forestry Science & Technology* (江苏林业科技), **34** (6): 23—27
- Wang Z (王忠), Dong SY (董仕勇), Luo YY (罗燕燕) *et al.*, 2008. Invasive plants in Guangzhou, China [J]. *Journal of Tropical and Subtropical Botany* (热带亚热带植物学报), **16** (1): 29—38
- Wei MY (韦美玉), Liu LP (刘丽萍), Wen ZR (文治瑞), 2006a. Growth and control of exotic plant in Qiannan area of Guizhou [J]. *Guizhou Agricultural Sciences* (贵州农业科学),

- 34 (2): 35—38
- Wei YL (韦原莲), Ye D (叶铎), Wen YG (温远光) *et al.*, 2006b. Invasive plant species in Shiwandashan Mountain National Natural Reserve, Guangxi, China [J]. *China Forestry Science and Technology* (林业科技开发), 20 (6): 23—26
- Wu T (吴彤), Meng C (孟陈), Dai J (戴洁) *et al.*, 2006. Exotic plants in Shandong Province [J]. *Journal of Shandong Normal University* (Natural Science) (山东师范大学学报(自然科学版)), 21 (4): 105—109
- Xie HY (谢红艳), Gong YZ (龚玉子), Zuo JF (左家哺), 2007a. A preliminary research of invasive plants in Nanyue Nature Reserve [J]. *Hunan Forestry Science & Technology* (湖南林业科技), 34 (2): 22—24
- Xie YZ (谢云珍), Wang YB (王玉兵), Tan WF (谭伟福), 2007b. Invasive plants in Guangxi Province [J]. *Journal of Tropical and Subtropical Botany* (热带亚热带植物学报), 15 (2): 160—167
- Xu CD (徐成东), Dong XD (董晓东), Lu SG (陆树刚), 2006. Invasive plants in Honghe River basin of Yunnan Province, China [J]. *Chinese Journal of Ecology* (生态学杂志), 13 (10): 57—58, 72.
- Xu CD (徐成东), Lu SG (陆树刚), 2006. The invasive plants in Yunnan [J]. *Guihaia* (广西植物), 26 (3): 227—234
- Xu HG, Qiang S, 2004. Inventory invasive alien species in China [J]. Beijing: China Environmental Science Press, 45—258
- Xu L (徐亮), Chen GX (陈功锡), Zhang DG (张代贵) *et al.*, 2009. Investigation on alien invasive plants in Xiangxi of Hunan Province, China [J]. *Journal of Jishou University* (Natural Science Edition) (吉首大学学报(自然科学版)), 30 (1): 98—103
- Yan YH (严岳鸿), He ZX (何祖霞), Gong Q (龚琴) *et al.*, 2007. The alien plant species in Guangzhou, China [J]. *Guihaia* (广西植物), 27 (4): 570—575
- Yan YH (严岳鸿), He ZX (何祖霞), Shea SS (余书生) *et al.*, 2005. A survey on the invasive plants in Kap O Islands, North-eastern Hong Kong [J]. *Bulletin of Botanical Research* (植物研究), 25 (2): 242—248
- Yan YH (严岳鸿), Xing FW (邢福武), Huang XX (黄向旭) *et al.*, 2004. Exotic plants in Shenzhen, China [J]. *Guihaia* (广西植物), 24 (3): 232—238
- Yang H (杨红), 2009. Present situation of invasive species in Qionghai wetland and their impact on its ecosystem [J]. *Journal of Mianyang Normal University* (绵阳师范学院学报), 28 (11): 58—62
- Yang J (杨坚), Chen HB (陈恒彬), 2009. A preliminary study on alien invasive plants in Fujian Province [J]. *Subtropical Plant Science* (亚热带植物科学), 38 (3): 47—52
- Yang JC (杨景成), Wang GM (王光美), Jiang CD (姜闯道) *et al.*, 2009. Ecological characters and distribution of invasive plants under the influence of urbanization in Beijing, China [J]. *Ecology and Environmental Sciences* (生态环境学报), 18 (5): 1857—1862
- Yang L (杨丽), Deng HP (邓洪平), Han M (韩敏) *et al.*, 2008. Pest risk analysis and assessment of invasive plants on environment in Chongqing [J]. *Journal of Southwest China Normal University* (Natural Science Edition) (西南师范大学学报(自然科学版)), 33 (1): 72—76
- Yi YJ (衣艳君), Li XS (李修善), Qiang S (强胜), 2005. Study on exotic weeds in Shandong Province [J]. *Territory & Natural Resources Study* (国土与自然资源研究), 3: 87—88
- Yu GL (于广丽), Li WZ (李文增), Chen GL (陈光磊), 2009. Ecological risk of invasive plants in Henan Province based on PRA [J]. *Journal of Henan Normal University* (Natural Science) (河南师范大学学报(自然科学版)), 37 (4): 133—135
- Zeng XF (曾宪锋), 2003. A preliminary study on 5 species invasive plants in E. Guangdong [J]. *Journal of Hanshan Teachers College* (韩山师范学院学报), 24 (3): 64, 69—71
- Zeng XF (曾宪锋), Lin XD (林晓单), Qiu HY (邱贺媛) *et al.*, 2009. Investigation on alien plants in E. Guangdong [J]. *Journal of Fujian Forestry Science and Technology* (福建林业科技), 36 (2): 174—179, 249
- Zhang JG (张建国), Zhang MR (张明如), 2009. Research on the establishment of the early-warning system of invasion of injurious biology outside Zhejiang [J]. *Journal of Fujian Forestry Science and Technology* (福建林业科技), 36 (1): 100—105
- Zhang XL (张绪良), Li YK (李永科), Xu ZJ (徐宗军) *et al.*, 2010. Research on invasion of harmful alien plants and their control strategies in Shandong Province [J]. *Hubei Agricultural Sciences* (湖北农业科学), 49 (1): 82—86
- Zhang ZX (张中信), 2009. Investigation on invasive plants in the new campus of Anqing normal University [J]. *Anhui Agricultural Science Bulletin* (安徽农学通报), 15 (20): 48—50
- Zhao H (赵宏), Dong CL (董翠玲), 2007. An investigation on the alien invasive plants in Kunyu mountain, Shandong Province [J]. *Jiangxi Science* (江西科学), 25 (4): 390—396
- Zhao JL (赵金丽), Ma YX (马友鑫), Zhu H (朱华) *et al.*, 2008. Invasion patterns of seven alien plant species along roadsides in southern mountainous areas of Yunnan Province [J]. *Biodiversity Science* (生物多样性), 16 (4): 369—380
- Zhao JM (赵见明), 2007. Study on the main exotic invasive plant species in Ruili, Yunnan [J]. *Journal of Southwest Forestry College* (西南林学院学报), 27 (1): 20—24
- Zhou XG (周小刚), Chen QH (陈庆华), Zhang H (张辉) *et al.*, 2008. Invasive alien weeds species in farmland and forest in Sichuan Province [J]. *Southwest China Journal of Agricultural Sciences* (西南农业学报), 21 (3): 852—858
- Zhu CS (朱长山), Tian CY (田朝阳), Lü SF (吕书凡) *et al.*, 2007. Investigation on and statistical analysis of alien invasive plants in Henan Province [J]. *Journal of Henan Agricultural University* (河南农业大学学报), 41 (2): 183—187